

Impact  
Factor  
2.147

ISSN 2349-638x

Refereed And Indexed Journal



**AAYUSHI  
INTERNATIONAL  
INTERDISCIPLINARY  
RESEARCH JOURNAL  
(AIIRJ)**

Monthly Publish Journal

VOL-III

ISSUE-XI

Nov.

2016

Address

- Vikram Nagar, Boudhi Chouk, Latur.
- Tq. Latur, Dis. Latur 413512
- (+91) 9922455749, (+91) 9158387437

Email

- aiirjpramod@gmail.com

Website

- www.aiirjournal.com

CHIEF EDITOR – PRAMOD PRAKASHRAO TANDALE

**Spatial Pattern Of Crop Combination In Kolhapur District (Maharashtra)**

**Shashikant Patil**

Asst. Professor,  
Mahavir Mahavidyalaya, Kolhapur.

**Dr. Arun Patil**

Head, Department of Geography,  
Mahavir Mahavidyalaya, Kolhapur

**Introduction**

India is an essentially agricultural country. In developing country like India crops generally grown in combination. The study of crop combination regions constitutes an important aspect of agricultural, geography and it provides a good basis for agricultural regionalization for a comprehensive and better understanding of the agricultural mosaic of an agro-climatic region. The study of crop combination is of greater significance even planning and development of agriculture a systematic crop combination has engaged attention of geographers and agriculture land use planners (Hussain M. 2002). The present study of crop combination analysis is related to the Kolhapur district which is part of Deccan plateau. Kolhapur district is one of agricultural most developed district of Maharashtra.

**Study region**


Kolhapur district is located in the south most part of Maharashtra Kolhapur district is situated in the southwestern part of Maharashtra. It lies between 15° 43' North to 17° 10' North latitude and 73°40' East to 74° 42' East longitude. Total area of Kolhapur district is 7692 Sq.km which occupies 2.62% area of total area of Maharashtra state. Kolhapur district comprising the Valleys of Warna, Panchaganga and their tributaries has a fertile & productive land.

• **Physiography and Climate**

The transitional geographical location of the district between Konkan coastal low land to the west and Deccan plateau to the east presents a variety in the geographical environment. General slop of the district is towards east and south-east. The general altitude of district of 1000 mts. to the west and 600 mst. to the east. The district has two main physiographic division's i.e. western hilly region and western hilly region consist of Phanhala, Shahuwadi, Gaganbawada, Radhanagari, Bhudargad, Aajra and Chandgad tahsils. The eastern plain region includes Shirol, Hatkanangale, Karvir, Kagal tahsils. The study region has developed drainage pattern. The rivers like Pachaganga, Varana, Dudhganga, Vedhganga, Hiranyakeshi and their tributaries play an important role in the development of agricultural in the study region.

The Kolhapur district has temperate climate. It receives rainfall mainly from south-west monsoon and intensity of rainfall decrease from west to east. The mean

temperature of the district lies between  $40^{\circ}\text{C}$  to  $16^{\circ}\text{C}$  in winter months. It exceeds more than  $38^{\circ}\text{C}$  in summer especially in April.



### Objectives of the study

1. To delineate crop combination regions of Kolhapur district.
2. To analyze the spatial pattern of crop combinations.

### Data source and Methodology

Present research work is based on secondary data . Data is collected from the following sources Socio economic review of Kolhapur District (2014-15), Agricultural Department of Kolhapur District.

Here an attempts has been made to study tahsil wise crop combination by maximum positive daviation method of Raffiullah . It is analyzed with the help of map. The Technique devised by raffiullah is as follows:

$$d = \sqrt{\frac{\sum d^2 p - \sum d^2}{N^2}}$$

d = deviation between actual crop percentage and the appropriate crop percentage in the

theoretical distribution.

p = positive difference


n = is negative difference

N = the number of crops in the crop combination

For the delineation of crop combination tahsil is selected as basic unit. present crop combination is limited for year of 2014-15 and for selected 13 crops.

#### **Tahsilwise area under cultivation of different crops.**

Physiography, rainfall, soil, irrigation facilities determine the productivity of different crops. The variations in above factors directly impact on crop cultivation. Table no.1 shows the tahsilwise area under cultivation of different crops and their percentage with total cultivated area of the district. Which shows tahsilwise cropping pattern of the district.



**Table No.1**  
**Tahsilwise area under cultivation of different crops ( 2014-15)**

Tahsils	Rice	Wheat	Jawar	Bajara	Maize	Nagali	Other cereals	Pulses	peanuts	sun flower	Other oil seeds	Sugar cane	Cotton	Total
Shahuwadi	50.96	4.00	2.00	0.00	0.80	9.55	2.51	7.03	10.85	0.15	1.97	10.18	0	100
Panhala	36.26	1.20	6.34	0.00	0.77	4.98	0.68	8.12	7.56	0	2.43	31.64	0	100
Hatkanangale	1.79	1.91	11.88	0.04	1.14	0.10	0.38	9.35	14.74	0.06	34.07	24.89	0	100
Shirol	1.13	2.65	4.45	0.04	0.66	0.09	0.76	7.59	4.44	0.01	42.81	35.32	0.05	100
Karveer	24.65	1.74	3.82	0.00	0.89	1.34	1.24	6	12.83	0.16	5.9	41.42	0.01	100
Gaganbawada	23.17	0.48	0.00	0.00	4.13	7.68	1.18	1.96	3.2	0	0.66	57.53	0	100
Radhanagari	46.28	0.39	2.62	0.00	2.28	10.31	1.71	0.78	4.8	0.01	2.05	28.79	0	100
Kagal	21.67	2.48	7.65	0.00	1.54	0.74	1.25	10.85	25.26	0.28	10.41	17.86	0	100
Bhudargad	50.17	1.55	0.21	0.00	0.84	13.95	0.85	5.31	12.16	0	1.55	13.4	0	100
Ajara	32.83	4.56	2.07	0.00	2.63	13.40	1.18	9.93	17.61	0	7.16	8.56	0.07	100
Gadhinglaj	13.80	1.31	6.14	0.00	2.33	0.70	1.04	8.31	36.13	0.07	21.11	9.06	0	100
Chandgad	35.93	0.36	1.04	0.00	1.03	18.83	0.47	2.53	17.1	0	0.68	21.88	0.16	100
Kolhapur District	2.43	3.02	6.22	0.01	1.85	6.80	1.30	8.97	19.79	0.09	18.45	31.05	0.03	100

Source: computed by researcher Analysis of spatial pattern of crop Combination (2014-15)

It has noted from table no. 2 that out of 12 tahsils of the district 7 tahsils come under the category of monoculture from the remaining tahsils 3 tahsils have 2 crop combinations and 2 tahsils have 3 crop combinations these crop combinations are also shown in map no. 2

The details of crop combinations in Kolhapur district by maximum positive deviation method of Raffiullah is analyzed as under.

**Table No.2**  
**Tahsilwise crop combination in Kolhapur District (2014-15)**

Sr.No.	Tahsil	Index Value of Crop Combination	Category	Crops in combination
1	Shahuwadi	129.49	Three crop	Rice -Peanuts -sugarcane
2	Panahala	523.30	Three crop	Rice –Sugarcane- Pulses
3	Karveer	73.61	Monoculture	Sugar
4	Hatkanangale	253.76	Monoculture	Oil seed
5	Shirol	945.68	Three crop	Oil seed-sugarcane-pulses
6	Gaganbawada	263.70	Two Crop	Sugar -Rice
7	Radhanagari	109.61	Two Crop	Rice - Sugar
8	Kagal	612.06	Monoculture	Peanuts
9	Bhudargad	127.88	Two Crop	Rice - Nagali
10	Ajara	294.80	Monoculture	Rice
11	Gadhinglaj	192.37	Monoculture	Peanuts
12	Chandagad	197.96	Monoculture	Rice


**A. Monoculture**

**I. Rice :-** Rice as a monoculture observed in two tahsils of Kolhapur district. These are Chandagad and Ajara. They cover an area about 35.93% and 32.83% of total sown area of that tahsils. The variance of the rice as monoculture is 294.80 in Ajara, and 197.96 in Chandagad. Due to high intensity of rainfall these tahsil have monoculture as rice

**II. Sugarcane :-** Sugarcane has a monoculture observed in Karvir tahsil. Sugarcane accounts 41.42% of total cropped area of Karvir tahsil. The index value of sugarcane is 73.60%. The tahsil Karvir has moderate and regular rainfall. The tahsil has well developed irrigation facilities and black fertile soil leads to the production of sugarcane

**III. Oil Seed:-** In tahsil Hatkanangale oil seed is as a monoculture. Out of total cultivated area 34.07% of area is under the cultivation of oil seeds. The variance of oil seed is 253.76. Fertile black soil well developed irrigation facility leads to the cultivation of oil seeds.

**IV. Peanuts :-** The tahsil of Gadhinglaj and kagal come under monoculture of peanuts. Area under peanuts cultivation in Gadhinglaj tahsil is 36.13% and 25.26% in kagal tahsil. The variance of peanuts in Gadhinglaj is 192.37 and in kagal tahsil it is 612.06. Fertile brownish black soil and developed irrigation facilities favourable for the cultivation of peanuts.



Map No.2

## B. Two crop combination

**I. Sugarcane and Rice :-** Radhanagari and Gaganbawada tahsils have 2 crop combinations. Sugarcane and Rice are two major crop in this tahsils. Sugarcane cover 57.53% out of total cultivated area in Gaganbawada where as 28.79% of area covered in Radhanagari on other side 23.17% cultivated area under the cultivation of Rice in Gaganbawada and 46.28% of area in Radhanagari. The variance of these two crop combination is 263.70. High amount of rainfall, fertile soil and irrigation facility leads to the cultivation of Sugarcane and Rice.

**II. Rice and Nagali :-** Due to high intensity of rainfall, Rice and Nagali these two crops are widely cultivated in Bhudargad tahsil. Out of total cultivated area 50.17% of area under the cultivation of Rice and 13.95% area under the cultivation of Nagali. The index value of these two crop combination is 127.88.

**C. Three crop combination:-** The three crop combination is observed in the Shahuwadi Panhala and Shirol tahsils

### I. Rice, Peanuts and Sugarcane

In Shahuwadi tahsil Rice, Peanuts and Sugarcane are the major crops cultivated. The tahsil Shahuwadi is well-suited for three crop combination. The crops Rice, Peanuts and Sugarcane are the first, second and third ranking crops and they covers an area about 50.96% , 10.85% and 10.18% respectively. The variance of the combination is 129.49

The tahil has rainfall ranging between 150 cm. and 250cm. and sometimes it is more than this. The tahsil has the shallow coarse soil in the hilly region. Which is best suitable for peanuts. in the river valleys of the tahsil fertile soil is present which is most useful for the cultivation of Rice and Sugarcane. Tahsil has also well developed irrigation facilities. All these conditions are useful for the dominance of rice, peanuts and sugarcane.

**II. Rice, Sugarcane and Pulses :-** Panhala tahsil has also three crop combination. Rice as first ranking crop 36,26.% , Sugarcaane as second ranking crop 31.64% and Peanuts as a third ranking crop (7.56%). The index value of three crop combination for panhala tahsil is 523.30

In Panhala tahsil, intensity of rainfall decrease from west to east. Therefore rice become dominant crop in western part and sugarcane and pulses are cultivated in eastern part of tahsil. Well developed fertile soil in the river valley region and irrigation facilities favorable for the cultivation of which favors for the cultivation of Rice and Sugarcane.

**III. Oil seeds, Sugarcane and Pulses :-** The tahsil Shirol have three crop combinations. Oil seeds, sugarcane and pulses are dominant crops. Out of total cultivated area 42.81% of area is under cultivation of oil seeds 35.32% of area is under cultivation of sugarcane and 7.59% area under cultivation of pulses.

The reason for the cultivation of oil seeds, sugarcane and pulses is that in western part of the tahsil medium deep soil is present. In the eastern part of the tahsil coarse shallow soil of laterite origin is the dominant soil. The fertility of the soil is increased by floods from the rivers like Krishna and Panchaganga.

### **Conclusion**

The crop combination in study region is the direct impact of the rainfall soils and irrigation facilities prevail there. In Panhala, Gaganbawada, Kagal and Ajara tahsils rice as a monoculture crop is present. This is due to the high amount of rainfall. Peanut as a monoculture crop of the tahsil Gadhinglaj and this is the impact of *Halki Kali Mati* present there. The tahsil Hatkanangale has soyabean as a monoculture crop. It is caused by the decrease in the amount and intensity of rainfall. Due to the presence of fertile soil, well developed irrigation facilities and moderate rainfall, Karvir tahsil has sugarcane as a dominant and monoculture crop.

Shahuwadi, bhudargarh and Shirol tahsils have the two crop combination. Tahsil Shiro has the soyabean-sugarcane combination. This is caused due to the low amount and intensity of rainfall and presence of fertile soil and irrigation facilities. Shahuwadi and Bhudargarh tahsils have Rice Nagali combination and Rice-Peanut combination respectively, which is also an impact of highest amount of rainfall and shallow coarse and laterite type of soil present there.

In the tahsil Radhanagari, Rice-Sugarcane-Nagali combination is present whereas, Rice-Nagali-Sugarcane combination has been observed in the Chandgad tahsil. This is also an impact of high amount of rainfall and good quality soil.

It has been concluded that more than 50% tahsils, i.e. in the seven have rice as a first ranking crop. Other important crops include sugarcane, Nagali and Peanut, Soyabean is the first ranking crop in the eastern part of the study region (viz in Hatkanangale and Shiro tahsils) where amount and intensity of the rainfall is low.

### **References**

- Census of India, (1991): District Census Handbook of Kolhapur.  
Husain, M. (2002): Systematic Agricultural Geography, Rawat Publication, Jaipur.  
Kharip Hangam Gutwar Pikawar Perchhetra Antim Ahwal (2003-2004): Krushi Vibhag, illa Parishad, Kolhapur.  
Kuniyal J.C.(1998): Crop Combination Regions in District Naital U.P.Himalaya. The Deccan Geographer, Vol.XXVI No.1,p.271-281.  
www.indiangos.com.